

---

**CBSE SAMPLE PAPER-02**

**Class – XI**

**MATHEMATICS**

---

Time allowed: 3 hours, Maximum Marks: 100

**General Instructions:**

- a) All questions are compulsory.
  - b) The question paper consists of 26 questions divided into three sections A, B and C. Section A comprises of 6 questions of one mark each, Section B comprises of 13 questions of four marks each and Section C comprises of 7 questions of six marks each.
  - c) All questions in Section A are to be answered in one word, one sentence or as per the exact requirement of the question.
  - d) Use of calculators is not permitted.
- 

**Section A**

**1. Identify a function  $f(x)$  so that  $f(x) \cdot f(y) = f(x + y)$**

**Sol:**  $f(x) = a^x$

$$\begin{aligned} f(x) \cdot f(y) &= a^x \cdot a^y \\ &= a^{x+y} = f(x+y) \end{aligned}$$

**2. If  $A = \{(x, y) : y = ax, x \in \mathbf{R}\}$  and  $B = \{(x, y) : y = a-x, x \in \mathbf{R}\}$  then what is  $(A \cap B)$**

**Sol:** When  $x = 0, y = 1$  in both cases. Hence

$$(A \cap B) = \{0, 1\}$$

**3. If  $R$  is a relation from a set  $A$  containing  $p$  elements to a set  $B$  containing  $q$  elements the find the number of subsets of  $A \times B$**

**Sol:**  $2^{pq}$

**4. Check whether the given lines are parallel or perpendicular.**

$$ax - by + c = 0 \quad \text{and} \quad \frac{ax}{2} - \frac{by}{2} + d = 0$$

**Sol:** They are parallel since

---

$$\begin{vmatrix} a & -b \\ \frac{a}{2} & \frac{-b}{2} \end{vmatrix} = 0$$

---

5. Find the area of the triangle whose vertices are (2,0),(5,3),(2,6)

Sol: Area of a triangle

$$\frac{1}{2} \begin{vmatrix} 2-2 & 0-6 \\ 5-2 & 3-6 \end{vmatrix} = \frac{1}{2} \begin{vmatrix} 0 & -6 \\ 3 & -3 \end{vmatrix} \\ = 9$$

---

6. Write the equation of a circle with center (0,0) and radius 5.

Sol:  $x^2 + y^2 = 25$

---

Section B

7. Solve  $\cos 3x = -\frac{1}{2}$

Sol:

$$\cos 3x = \cos \frac{2\pi}{3}$$

$$3x = 2n\pi \pm \frac{2\pi}{3}$$

$$x = \frac{2n\pi}{3} \pm \frac{2\pi}{9}, n \in Z$$

---

8. Prove by mathematical induction that  $1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$ .

Sol: Let  $P(n)$  be the statement given by

$$1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

$$P(1) = \frac{1(1+1)}{2} = 1, \text{ True}$$

Let it be true for  $n=m$

$$1 + 2 + 3 + \dots + m = \frac{m(m+1)}{2}$$

$$1 + 2 + 3 + \dots + m + (m + 1) = \frac{m(m+1)}{2} + (m + 1)$$

---

$$P(m+1) = \frac{m(m+1)}{2} + (m+1)$$

$$P(m+1) = \frac{m^2+3m+2}{2}$$

$$P(m+1) = \frac{(m+1)(m+2)}{2}$$

Thus  $P(m)$  is true  $\Rightarrow P(m+1)$  is True

**9. Find the square root of  $\sqrt{-8i}$ .**

**Sol:** Let  $\sqrt{z} = \sqrt{-8i}$

$$\sqrt{z} = \pm \left\{ \frac{\sqrt{|z| - \operatorname{Re}(z)}}{\sqrt{2}} \right\} - i \left\{ \frac{\sqrt{|z| - \operatorname{Re}(z)}}{\sqrt{2}} \right\}, \operatorname{Im}(z) < 0$$

$$\sqrt{-8i} = \pm \left\{ \frac{\sqrt{8+0}}{\sqrt{2}} - i \frac{\sqrt{8-0}}{\sqrt{2}} \right\}, \operatorname{Im}(z) < 0$$

$$= \pm(2 - 2i)$$

**10. Solve the inequality  $\frac{2x+5}{x-2} \geq 3$ .**

**Sol:**  $\frac{2x+5}{x-2} - 3 \geq 0$

**11. Find the value of  $x$  if  $|2Cx| = |2Cx+4|$ .**

**Sol:**  $x + x + 4 = 12$

$$2x = 8$$

$$x = 4$$

**12. Three cars are there in a race. Car A is 3 times as likely to win as car B. Car B is twice as likely to win as car C. What is the probability of winning each car.**

**Sol:** Let  $p$  be the probability of winning Car C,

$$P(C)$$

$$P(C) = p$$

$$P(B) = 2p$$

$$P(A) = 6p$$

$$P(A) + P(B) + P(C) = 1$$

---

$$p + 2p + 6p = 1$$

$$9p = 1$$

$$p = \frac{1}{9}$$

$$P(C) = \frac{1}{9}$$

$$P(B) = \frac{2}{9}$$

$$P(A) = \frac{6}{9}$$

---

**13. If  $f(x)$  is a function that contains 3 in its domain and range and satisfy the relation  $f(f(x)).(1 + f(x)) = -f(x)$  find  $f(3)$**

**Sol:** Let  $a$  satisfy the relation  $f(a) = 3$

$$f(f(a)).(1 + f(a)) = -f(a)$$

$$f(3).(4) = -3$$

$$f(3) = -\frac{3}{4}$$

---

**14. If  $\tan A = \frac{1}{3}$  and  $\tan B = \frac{1}{2}$  prove that  $\sin 2(A + B) = 1$ .**

**Sol:**

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$= \frac{\frac{1}{3} + \frac{1}{2}}{1 - \frac{1}{3} \cdot \frac{1}{2}} = 1$$

$$A + B = 45$$

$$2(A + B) = 90$$

$$\sin 90 = 1$$

---

**15. Find two numbers such that their arithmetic mean is 15 and Geometric mean is 9 without using the identity  $(a + b)^2 = (a - b)^2 + 4ab$**

**Sol:** Form a quadratic equation sum of whose roots are 30 and product of the roots is 81

$$x^2 - x(30) + 81 = 0$$

$$x^2 - 3x - 27x + 81 = 0$$

$$x(x - 3) - 27(x - 3)$$

---

$$(x - 3)(x - 27) = 0$$

Hence the numbers are 3 and 27

---

**16. Let  $f : R \rightarrow R$  be a function given by  $f(x) = x^2 + 2$  find  $f^{-1}(27)$**

**Sol:** Let  $f : R \rightarrow R$  be a function given by  $f(x) = x^2 + 2$  find  $f^{-1}(27)$

$$f(x) = x^2 + 2$$

$$x^2 + 2 = 27$$

$$x^2 = 25$$

$$x = \pm 5$$

$$f^{-1}(27) = \{-5, 5\}$$

---

**17. Find the domain and range of the function  $f(x) = \frac{x-a}{a+1-x}$  where  $a$  is a positive integer.**

**Sol:** The function is defined for all values of  $x$  where the denominator is not equal to zero

$$a + 1 - x \neq 0$$

$$\text{Hence domain} = R - \{(a + 1)\}$$

Range of  $f$

$$\text{Let } y = f(x)$$

$$y = \frac{x-a}{a+1-x}$$

$$(a+1)y - xy = x-a$$

$$x(y+1) = (a+1)y + a$$

$$x = \frac{(a+1)y + a}{y+1}$$

$$\text{Range of } f = R - \{-1\}$$

---

**18. Find the limit of  $\lim_{x \rightarrow 0} \frac{\sqrt{a+x} - \sqrt{a}}{x}$**

**Sol:** Rationalize the numerator

$$\begin{aligned} & \lim_{x \rightarrow 0} \frac{\sqrt{a+x} - \sqrt{a}}{x} \\ &= \lim_{x \rightarrow 0} \frac{(\sqrt{a+x} - \sqrt{a})(\sqrt{a+x} + \sqrt{a})}{x(\sqrt{a+x} + \sqrt{a})} \\ &= \lim_{x \rightarrow 0} \frac{x}{x(\sqrt{a+x} + \sqrt{a})} \\ &= \frac{1}{2\sqrt{a}} \end{aligned}$$

19. Find the sign and value of the expression  $\sin 75^\circ + \cos 75^\circ$

Sol:

$$\begin{aligned} & \sin 75^\circ + \cos 75^\circ \\ &= \sqrt{2} \left( \frac{1}{\sqrt{2}} \sin 75^\circ + \frac{1}{\sqrt{2}} \cos 75^\circ \right) \\ &= \sqrt{2} (\cos 45^\circ \sin 75^\circ + \sin 45^\circ \cos 75^\circ) \\ &= \sqrt{2} \sin(75^\circ + 45^\circ) \\ &= \sqrt{2} \sin 120^\circ \end{aligned}$$

Hence sign is positive and value is  $\frac{\sqrt{2} \cdot \sqrt{3}}{2} = \frac{\sqrt{6}}{2}$

### Section C

20. In how many ways can 3 students from Class 12, 4 from class 11, 4 from class 10 and 2 from class 9 be seated in a row so that those of the same classes sit together. Also find the number of ways they can be arranged in at a round table.

Sol: There are 4 groups and four groups can be arranged in  $4!$  ways. Class 12 can be arranged in  $3!$  ways, Class 11 can be arranged in  $4!$  Class 10 can be arranged in  $4!$ . Class 9 can be arranged in ways. Hence Total number of ways that they can be arranged in a row  $4! \times 3! \times 4! \times 4! \times 2! = 165888$

In a circular seating arrangement the four groups can be arranged only in  $3!$  ways only.

---

Hence the total number of ways that they can be seated at a round table =  $3! \times 3! \times 4! \times 4! \times 2! = 41472$

---

**21. A circle represented by the equation  $(x - a)^2 + (y - b)^2 = r^2$**

**This makes two complete revolutions along the positive direction of the x-axis. Find the equation of the circle in the new position.**

**Sol:** The new coordinates of the centre in the new position are

$$(a + 4\pi r, b)$$

$$\{x - (a + 4\pi r)\}^2 + (y - b)^2 = r^2$$

---

**22. Show that the equation  $x^2 + 4y^2 + 4x + 16y + 16 = 0$  represents an ellipse.**

**Sol:**

$$x^2 + 4y^2 + 4x + 16y + 16 = 0$$

$$x^2 + 4x + 4 + 4y^2 + 16y + 16 = 4$$

$$(x + 2)^2 + 4(y + 2)^2 = 4$$

$$\frac{(x + 2)^2}{2^2} + \frac{(y + 2)^2}{1^2} = 1$$

This equation represents an ellipse.

---

**23. Calculate the mean deviation about the mean from the following data**

xi	2	15	17	23	27
fi	12	6	12	9	5

**Sol:**

$x_i$	$f_i$	$f_i x_i$	$ x_i - 15 $	$f_i  x_i - 15 $
2	12	24	13	156
15	6	90	0	0
17	12	204	2	24
23	9	207	8	72
27	5	135	12	60
	$N = \sum f_i = 44$	$\sum f_i x_i = 660$		$f_i \sum  x_i - 15  = 312$

$$\text{Mean} = \bar{X} = \frac{1}{N} (\sum f_i x_i)$$

$$= \frac{660}{44} = 15$$

$$\text{Mean Deviation} = \text{M.D} = \frac{1}{N} (\sum f_i |x_i - 15|)$$

$$= \frac{312}{44} = 7.0909$$

**24. If the ratio of the roots of the equation  $x^2 + px + q = 0$  is the same as  $x^2 + p_1 x + q_1 = 0$  then prove that  $p^2 q_1 = p_1^2 q$**

**Sol:** Let the ratios be  $a : b$

$$x^2 + px + q = 0$$

$$a\alpha + b\beta = -p$$

$$a\beta + b\alpha = -p_1$$

$$a\alpha \times b\alpha = q$$

$$a\beta \times b\beta = q_1$$

$$(a + b)\alpha = -p$$

$$(a + b)\beta = -p_1$$

$$ab\alpha^2 = q$$

$$ab\beta^2 = q_1$$

$$\frac{(a+b)^2 \alpha^2}{(a+b)^2 \beta^2} = \frac{p^2}{p_1^2}$$

$$\frac{\alpha^2}{\beta^2} = \frac{p^2}{p_1^2}$$

$$\frac{\alpha^2}{\beta^2} = \frac{q}{q_1}$$

$$\frac{p^2}{p_1^2} = \frac{q}{q_1}$$

$$p^2 q_1 = p_1^2 q$$

25. Prove that  $a \cdot a^{\frac{1}{2}} \cdot a^{\frac{1}{4}} \cdot a^{\frac{1}{8}} \dots \infty = a^2$ .

Sol:

$$1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots \infty$$

$$= \frac{1}{1 - \frac{1}{2}} = \frac{1}{\frac{1}{2}} = 2$$

$$a \cdot a^{\frac{1}{2}} \cdot a^{\frac{1}{4}} \cdot a^{\frac{1}{8}} \dots \infty = a^2$$

26. In a survey of 700 students in a medical college 200 went for regular entrance coaching, 295 attended only correspondence coaching, 115 attended both regular and correspondence coaching. Find how many got admission without any entrance coaching.

Sol: It is given that

$$n(U) = 700, n(A) = 200, n(B) = 295, n(A \cap B) = 115$$

We need to find out

$$n(A' \cap B')$$

$$n(A' \cap B') = n(A \cup B)'$$

$$= n(U) - n(A \cup B)$$

$$= n(U) - \{n(A) + n(B) - n(A \cap B)\}$$

$$= 700 - \{200 + 295 - 115\}$$

$$= 320$$